Syritta diversity in the Afrotropical Region

Kurt Jordaens^{*1}, Georg Goergen², John Midgley^{3,4}, Terence Bellingan^{5,6}, Bonolo Mosime^{7,8}, Marc De Meyer¹, and Jeroen Van Steenis⁹

³Department of Zoology Entomology, Rhodes University, Makhanda, 6139 – South Africa
⁴KwaZulu-Natal Museum, Pietermaritzburg, KwaZulu-Natal, 3021 – South Africa
⁵Department of Zoology Entomology, Rhodes University, Makhanda, 6139 – South Africa
⁶Department of Entomology Arachnology, Albany Museum, Makhanda, 6139 – South Africa
⁷KwaZulu-Natal Museum, Pietermaritzburg, KwaZulu-Natal, 3021 – South Africa
⁸University of KwaZulu-Natal, Pietermaritzburg, KwaZulu-Natal, 3021 – South Africa
⁹Naturalis Biodiversity Center [Leiden] – Netherlands

Abstract

The revision of the hoverfly genus Syritta Le Peletier & Serville, 1828 by Lyneborg & Barkemeyer (2005) comprises approximately 60 species worldwide, with 40 of these recorded from the Afrotropical Region. Over the last decade, we have collected and studied > 2,000 Syritta specimens from almost 20 Afrotropical countries and have used morphology and DNA barcoding to identify the species, and to link males to females. This has resulted in at least six new species and the discovery of the missing sex for eight species where only one of the sexes was known. Adding museum collections to the newly collected material allowed us to study the species distributions, diversity, biogeography and predict future distributions using modelling techniques. Moreover, an updated identification key will be provided including high-resulution stacking photographs for both males and females.

Keywords: South Africa, Biodiversity hotspots, biogeography, species distribution modelling

¹Royal Museum for Central Africa, Invertebrates Section, Leuvensesteenweg 13, 3080 Tervuren – Belgium

²International Institute of Tropical Agriculture (IITA), Biodiversity Centre, 08 BP 0932 Tri Postal, Cotonou – Benin

^{*}Speaker