

Please mark your choices with an "X".

Presentation:

Poster Presentation.....X

Oral Presentation.....

**Preimaginal morphology of the European endemic species *Merodon triangulum* Vujić, Radenković & Hurkmans, 2020 of the *Merodon constans* group (Diptera, Syrphidae).**

Andrea Aracil <sup>1</sup>, Andrijana Andrić <sup>2</sup>, Celeste Pérez-Bañón <sup>1</sup>, Snežana Radenković <sup>3</sup>, Ante Vujić <sup>3</sup> and Santos Rojo <sup>1</sup>

1 Department of Environmental Sciences & Natural Resources, University of Alicante, PO Box. 99, E-03080 Alicante, Spain.

E-mail: andrea.aracil@ua.es, celeste.perez@ua.es, santos.rojo@ua.es

2 University of Novi Sad, BioSense Institute, Dr Zorana Đinđića 1, 21000 Novi Sad, Serbia; e-mail: andrijana.andric@biosens.uns.ac.rs

3 Department of Biology and Ecology, University of Novi Sad, Trg Dositeja Obradovića 2, 21000 Novi Sad, Serbia. E-mail: snezana.radenkovic@dbe.uns.ac.rs, ante.vujic@dbe.uns.ac.rs

Keywords (up to 5): *Merodon*, Serbia, preimaginal morphology, SEM

The genus *Merodon* Meigen, 1803 is one of the richest hoverfly genera, with around 180 species known species in the Palearctic region, with the highest species diversity recorded for the Mediterranean Basin. The larval biology of most *Merodon* species remain unknown, however, the development of all species which the preimaginal stages have been identified, occurs in bulbs of monocotyledonous plants of families Amaryllidaceae, Asparagaceae, Iridaceae and Liliaceae.

The genus is classified into more than 20 monophyletic species (Vujić et al., 2021) groups being *Merodon triangulum* Vujić, Radenković & Hurkmans, 2020, belongs to the *constans* group and the *albifrons* lineage. It is a European endemic species, known from north-eastern Italy, northern Austria, eastern Hungary, Slovenia, Croatia, Serbia, Montenegro, and North Macedonia. The aim of this paper is to present the first data about preimaginal morphology (both larva and pupa) of this species, described using both optical ~~microscopy~~ and scanning electron microscopy. The larval description was obtained from wild larval specimens collected in Petrovaradinski rid, Novi Sad, Serbia, collected in June 2020, feeding inside bulbs of *Leucojum vernum* L. 1753 (Amaryllidaceae).

References:

Vujić A, Radenković S, Likov L, Veselić S (2021) Taxonomic complexity in the genus *Merodon* Meigen, 1803 (Diptera, Syrphidae). ZooKeys 1031: 85-124. Doi: 10.3897/zookeys.1031.62125